

EDINBURGH POSTNATAL DEPRESSION SCALE (EPDS)

The EPDS was developed for screening postpartum women in outpatient, home visiting settings, or at the 6–8 week postpartum examination. It has been utilized among numerous populations including U.S. women and Spanish speaking women in other countries. The EPDS consists of 10 questions. The test can usually be completed in less than 5 minutes. Responses are scored 0, 1, 2, or 3 according to increased severity of the symptom. Items marked with an asterisk (*) are reverse scored (i.e., 3, 2, 1, and 0). The total score is determined by adding together the scores for each of the 10 items. Validation studies have utilized various threshold scores in determining which women were positive and in need of referral. Cut-off scores ranged from 9 to 13 points. Therefore, to err on safety's side, a woman scoring 9 or more points or indicating any suicidal ideation – that is she scores 1 or higher on question #10 – should be referred immediately for follow-up. Even if a woman scores less than 9, if the clinician feels the client is suffering from depression, an appropriate referral should be made. The EPDS is only a screening tool. It does not diagnose depression – that is done by appropriately licensed health care personnel. Users may reproduce the scale without permission providing the copyright is respected by quoting the names of the authors, title and the source of the paper in all reproduced copies.

Instructions for Users

1. The mother is asked to underline 1 of 4 possible responses that comes the closest to how she has been feeling the previous 7 days.
2. All 10 items must be completed.
3. Care should be taken to avoid the possibility of the mother discussing her answers with others.
4. The mother should complete the scale herself, unless she has limited English or has difficulty with reading.

Name:

Date:

Address:

Baby's Age:

As you have recently had a baby, we would like to know how you are feeling. Please UNDERLINE the answer which comes closest to how you have felt **IN THE PAST 7 DAYS**, not just how you feel today.

Here is an example, already completed.

I have felt happy:

Yes, all the time

Yes, most of the time

No, not very often

No, not at all

This would mean: "I have felt happy most of the time" during the past week. Please complete the other questions in the same way.

In the past 7 days:

- | | |
|---|--|
| 1. I have been able to laugh and see the funny side of things
As much as I always could
Not quite so much now
Definitely not so much now
Not at all | *6. Things have been getting on top of me
Yes, most of the time I haven't been able to cope at all
Yes, sometimes I haven't been coping as well as usual
No, most of the time I have coped quite well
No, have been coping as well as ever |
| 2. I have looked forward with enjoyment to things
As much as I ever did
Rather less than I used to
Definitely less than I used to
Hardly at all | *7. I have been so unhappy that I have had difficulty sleeping
Yes, most of the time
Yes, sometimes
Not very often
No, not at all |
| *3. I have blamed myself unnecessarily when things went wrong
Yes, most of the time
Yes, some of the time
Not very often
No, never | *8. I have felt sad or miserable
Yes, most of the time
Yes, quite often
Not very often
No, not at all |
| 4. I have been anxious or worried for no good reason
No, not at all
Hardly ever
Yes, sometimes
Yes, very often | *9. I have been so unhappy that I have been crying
Yes, most of the time
Yes, quite often
Only occasionally
No, never |
| *5. I have felt scared or panicky for no very good reason
Yes, quite a lot
Yes, sometimes
No, not much
No, not at all | *10. The thought of harming myself has occurred to me
Yes, quite often
Sometimes
Hardly ever
Never |

EDINBURGH POSTNATAL DEPRESSION SCALE (EPDS)

J. L. Cox, J.M. Holden, R. Sagovsky

From: *British Journal of Psychiatry* (1987), 150, 782-786.

Edinburgh Postnatal Depression Scale¹ (EPDS)

Name: _____

Address: _____

Your Date of Birth: _____

Baby's Date of Birth: _____

Phone: _____

As you are pregnant or have recently had a baby, we would like to know how you are feeling. Please check the answer that comes closest to how you have felt **IN THE PAST 7 DAYS**, not just how you feel today.

Here is an example, already completed.

I have felt happy:

- Yes, all the time
- Yes, most of the time This would mean: "I have felt happy most of the time" during the past week.
- No, not very often Please complete the other questions in the same way.
- No, not at all

In the past 7 days:

- | | |
|--|--|
| <p>1. I have been able to laugh and see the funny side of things</p> <ul style="list-style-type: none"><input type="checkbox"/> As much as I always could<input type="checkbox"/> Not quite so much now<input type="checkbox"/> Definitely not so much now<input type="checkbox"/> Not at all <p>2. I have looked forward with enjoyment to things</p> <ul style="list-style-type: none"><input type="checkbox"/> As much as I ever did<input type="checkbox"/> Rather less than I used to<input type="checkbox"/> Definitely less than I used to<input type="checkbox"/> Hardly at all <p>*3. I have blamed myself unnecessarily when things went wrong</p> <ul style="list-style-type: none"><input type="checkbox"/> Yes, most of the time<input type="checkbox"/> Yes, some of the time<input type="checkbox"/> Not very often<input type="checkbox"/> No, never <p>4. I have been anxious or worried for no good reason</p> <ul style="list-style-type: none"><input type="checkbox"/> No, not at all<input type="checkbox"/> Hardly ever<input type="checkbox"/> Yes, sometimes<input type="checkbox"/> Yes, very often <p>*5. I have felt scared or panicky for no very good reason</p> <ul style="list-style-type: none"><input type="checkbox"/> Yes, quite a lot<input type="checkbox"/> Yes, sometimes<input type="checkbox"/> No, not much<input type="checkbox"/> No, not at all | <p>*6. Things have been getting on top of me</p> <ul style="list-style-type: none"><input type="checkbox"/> Yes, most of the time I haven't been able to cope at all<input type="checkbox"/> Yes, sometimes I haven't been coping as well as usual<input type="checkbox"/> No, most of the time I have coped quite well<input type="checkbox"/> No, I have been coping as well as ever <p>*7. I have been so unhappy that I have had difficulty sleeping</p> <ul style="list-style-type: none"><input type="checkbox"/> Yes, most of the time<input type="checkbox"/> Yes, sometimes<input type="checkbox"/> Not very often<input type="checkbox"/> No, not at all <p>*8. I have felt sad or miserable</p> <ul style="list-style-type: none"><input type="checkbox"/> Yes, most of the time<input type="checkbox"/> Yes, quite often<input type="checkbox"/> Not very often<input type="checkbox"/> No, not at all <p>*9. I have been so unhappy that I have been crying</p> <ul style="list-style-type: none"><input type="checkbox"/> Yes, most of the time<input type="checkbox"/> Yes, quite often<input type="checkbox"/> Only occasionally<input type="checkbox"/> No, never <p>*10. The thought of harming myself has occurred to me</p> <ul style="list-style-type: none"><input type="checkbox"/> Yes, quite often<input type="checkbox"/> Sometimes<input type="checkbox"/> Hardly ever<input type="checkbox"/> Never |
|--|--|

Administered/Reviewed by _____ Date _____

¹Source: Cox, J.L., Holden, J.M., and Sagovsky, R. 1987. Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry* 150:782-786 .

²Source: K. L. Wisner, B. L. Parry, C. M. Piontek, Postpartum Depression N Engl J Med vol. 347, No 3, July 18, 2002, 194-199

Users may reproduce the scale without further permission providing they respect copyright by quoting the names of the authors, the title and the source of the paper in all reproduced copies.

Edinburgh Postnatal Depression Scale¹ (EPDS)

Postpartum depression is the most common complication of childbearing.² The 10-question Edinburgh Postnatal Depression Scale (EPDS) is a valuable and efficient way of identifying patients at risk for "perinatal" depression. The EPDS is easy to administer and has proven to be an effective screening tool.

Mothers who score above 13 are likely to be suffering from a depressive illness of varying severity. The EPDS score should not override clinical judgment. A careful clinical assessment should be carried out to confirm the diagnosis. The scale indicates how the mother has felt **during the previous week**. In doubtful cases it may be useful to repeat the tool after 2 weeks. The scale will not detect mothers with anxiety neuroses, phobias or personality disorders.

Women with postpartum depression need not feel alone. They may find useful information on the web sites of the National Women's Health Information Center <www.4women.gov> and from groups such as Postpartum Support International <www.chss.iup.edu/postpartum> and Depression after Delivery <www.depressionafterdelivery.com>.

SCORING

QUESTIONS 1, 2, & 4 (without an *)

Are scored 0, 1, 2 or 3 with top box scored as 0 and the bottom box scored as 3.

QUESTIONS 3, 5-10 (marked with an *)

Are reverse scored, with the top box scored as a 3 and the bottom box scored as 0.

Maximum score: 30
Possible Depression: 10 or greater
Always look at item 10 (suicidal thoughts)

Users may reproduce the scale without further permission, providing they respect copyright by quoting the names of the authors, the title, and the source of the paper in all reproduced copies.

Instructions for using the Edinburgh Postnatal Depression Scale:

1. The mother is asked to check the response that comes closest to how she has been feeling in the previous 7 days.
2. All the items must be completed.
3. Care should be taken to avoid the possibility of the mother discussing her answers with others. (Answers come from the mother or pregnant woman.)
4. The mother should complete the scale herself, unless she has limited English or has difficulty with reading.

¹Source: Cox, J.L., Holden, J.M., and Sagovsky, R. 1987. Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry* 150:782-786.

²Source: K. L. Wisner, B. L. Parry, C. M. Piontek, Postpartum Depression N Engl J Med vol. 347, No 3, July 18, 2002, 194-199